

Amendments to the Claims

This listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of Claims:

1. (Currently amended) A process for producing 2-O- α -glucopyranosyl-L-ascorbic acid, comprising the steps of:

allowing α -isomaltosyl glucosaccharide-forming enzyme together with or without cyclomaltodextrin glucanotransferase (EC 2.4.1.19) to act on a solution comprising L-ascorbic acid and α -glucosyl saccharide selected from the group consisting of maltooligosaccharide, maltodextrin, cyclodextrin, amylose, amylopectin, soluble starch, liquefied starch, gelatinized starch, and glycogen to obtain a reaction mixture containing 2-O- α -glucopyranosyl-L-ascorbic acid in an amount of 10% (w/w) or higher wherein the reaction mixture also contains ~~and~~ each of 5-O- α -glucopyranosyl-L-ascorbic acid and 6-O- α -glucopyranosyl-L-ascorbic acid in an

amount of less than 0.1% (w/w), on a dry solid basis;
and

collecting the 2-O- α -glucopyranosyl-L-
ascorbic acid from the reaction mixture;
wherein said α -isomaltosyl glucosaccharide-
forming enzyme has an activity of forming a
saccharide with a glucose polymerization degree of 3
or higher and bearing both the α -1,6 glucosidic
linkage as a linkage at the non-reducing end and the
 α -1,4 glucosidic linkage other than the linkage at
the non-reducing end from a saccharide with a glucose
polymerization degree of 2 or higher and bearing the
 α -1,4 glucosidic linkage as a linkage at the non-
reducing end by α -glucosyl-transferring reaction
without increasing the reducing power of the reaction
mixture; wherein said α -isomaltosyl glucosaccharide-
forming enzyme is obtained from the genusgenera
Arthrobacter and Bacillus.

2. (Previously presented) The process of claim 1, wherein glucoamylase (EC 3.2.1.3) is allowed to act on the reaction mixture after the action of α -isomaltosyl glucosaccharide-forming enzyme on said solution together with or without cyclomalodextrin glucanotransferase.

Claims 3-5. (Cancelled)

6. (Previously presented) The process of claim 1, wherein the step of collecting 2-O- α -glucopyranosyl-L-ascorbic acid comprises a step of using a strongly-acidic cation exchange resin.

7. (Previously amended) The process of claim 1, wherein the formed 2-O- α -glucopyranosyl-L-ascorbic acid is collected in the form of a syrup, a powder, or a crystal.

Claims 8-20. (Cancelled)

21. (Previously presented) The process of claim 6, further comprising pulverizing or crystallizing the 2-O- α -glucopyranosyl-L-ascorbic acid.

Claims 22-23. (Cancelled)